

=>

Uploading C:\Program Files\Stnexp\Queries\10581322-broad.str



chain nodes :

14 15 17

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12

chain bonds :

6-14 8-17 14-15 15-17

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12

exact/norm bonds :

6-14 8-17 14-15 15-17

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12

isolated ring systems :

containing 1 :

G1:O,S

Connectivity :

15:2 E exact RC ring/chain

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom

11:Atom 12:Atom 14:Atom 15:CLASS 17:CLASS

Generic attributes :

14:

Saturation : Unsaturated

Number of Carbon Atoms : less than 7

Number of Hetero Atoms : 2 or more

Type of Ring System : Monocyclic

15:

Saturation : Saturated

Element Count :

Node 14: Limited

O,O1

N,N1

C,C3

=>
Uploading C:\Program Files\Stnexp\Queries\10581322-elected-species-Final.str

L4 STRUCTURE UPLOADED

=> d his

(FILE 'HOME' ENTERED AT 11:51:14 ON 16 JAN 2008)

FILE 'REGISTRY' ENTERED AT 11:51:24 ON 16 JAN 2008
ACT BRD581322/A

L1 STR
L2 (210227)SEA FILE=REGISTRY ABB=ON PLU=ON NOC3/ES
L3 2449 SEA FILE=REGISTRY SUB=L2 SSS FUL L1

L4 STRUCTURE UPLOADED
L5 21 S L4 SAM SUB=L3
L6 460 S L4 SSS FULL SUB=L3


FILE 'CAPLUS' ENTERED AT 11:52:02 ON 16 JAN 2008
L7 26 S L6
L8 2 S US200!-581322/APPS
L9 1 S L7 AND L8
L10 25 S L7 NOT L8

FILE 'REGISTRY' ENTERED AT 11:52:25 ON 16 JAN 2008

=> d l4



L4 HAS NO ANSWERS

L4 STR



G1 C,O,S

Structure attributes must be viewed using STN Express query preparation.



chain nodes :

8 10 12 19 20 21 22

ring nodes :

1 2 3 4 5 6 13 14 15 16 17 18

chain bonds :

5-8 8-10 10-12 12-15 19-20 20-21 20-22

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 13-14 13-18 14-15 15-16 16-17 17-18

exact/norm bonds :

5-8 8-10 10-12 12-15 19-20 20-21 20-22

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 13-14 13-18 14-15 15-16 16-17 17-18

isolated ring systems :

containing 1 : 13 :

G1:C,O,S

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 8:Atom 10:CLASS 12:CLASS 13:Atom
14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:CLASS 20:CLASS 21:CLASS 22:CLASS
23:Atom

Generic attributes :

8:

Saturation : Unsaturated

Number of Carbon Atoms : less than 7

Number of Hetero Atoms : 2 or more

Type of Ring System : Monocyclic

Element Count :

Node 8: Limited

O,O1



N,N1

C,C3

L10 ANSWER 1 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
 AN 2007:1089015 CAPLUS Full-text
 DN 147:416507
 TI Holographic recording material, recording method, and optical recording medium
 IN Nomura, Tomoko; Yamashita, Noriko; Takizawa, Hiroo
 PA Fuji Photo Film Co., Ltd., Japan
 SO Jpn. Kokai Tokkyo Koho, 59pp.
 CODEN: JKXXAF
 DT Patent
 LA Japanese
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI JP 2007248517	A	20070927	JP 2006-67903	20060313
PRAI JP 2006-67903		20060313		

GI





AB Title recording material includes a sensitizing dye, a dye precursor, and a basic compound selected from I and II (R1 = H, alkyl, alkenyl, cycloalkyl, aryl; 21-3 = substituent; a = 0-5; k = 1-10; m = 0-2; R2, R3 = H, alkyl, alkenyl, cycloalkyl; b, c = 0-3). The sensitizing dye absorbs the light for holog. exposure to become excited state. The dye precursor receives charge transfer or energy transfer from the excited sensitizing dye and decomps. to provide the dye in the presence of the basic compound

IT 950595-20-4
 RI: TEM (Technical or engineered material use); USES (Uses)
 (holog. recording material and recording method)

RN 950595-30-4 CAPLUS

CN 2-Propenoic acid, 2-cyano-3-[3,5-dichloro-4-[(2-{4-[(diethylamino)sulfonyl]-2-nitrophenyl]-2,3-dihydro-5-methyl-3-oxo-4-isoxazolyl)methoxy]phenyl]-, 2-hexyldecyl ester (CA INDEX NAME)

IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MN, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA
 PRAI US 2005-715214P P 20050907
 US 2006-789387P P 20060405
 OS MARPAT 146:337575
 GI





AB Comps. of formula I that are active on at least one of PPAR α , PPAR δ , and PPAR γ , which are useful for therapeutic and/or prophylactic methods involving modulation of at least one of PPAR α , PPAR δ , and PPAR γ , are described. Comps. of formula I wherein X is CO $_2$ H or deriv., CONH $_2$ and deriv., and carboxylic acid isostere; W is bond, (un)substituted C1-2 alkylamino, (un)substituted C1-2 alkoxy, (un)substituted C1-3 alkylene, (un)substituted ethylene, R1 and R2 are independently H, halo, (un)substituted lower alkyl, (un)substituted lower alkenyl, (un)substituted lower alkynyl, etc.; R3 is (un)substituted alkyl; L is O, S, NH and deriv., CO, CS, SO, SO $_2$, CONH and deriv., etc.; and their pharmaceutically acceptable salts, prodrugs, tautomers, and isomers, thereof are claimed. Example compound II was prepared by a multistep procedure (procedure given). All the invention comps. were evaluated for their PPAR α , PPAR δ , and PPAR γ activity. From the assay, it was determined that several comps. exhibited EC $_{50}$ values of less than or equal to 1 μ M against at least one of the PPAR α , PPAR δ , or PPAR γ .

IT 929092-14-4P

RI: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug candidate; preparation of arylacetic acids and related comps. useful in prophylaxis and treatment of diseases - mediated by PPAR α , PPAR δ and PPAR γ receptors)

RN 929092-94-4 CAPLUS


CN Benzenecarboxylic acid, 3-butoxy-5-[(3-{(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl)methoxy]- (CA INDEX NAME)



L10 ANSWER 2 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
 AN 2007:284433 CAPLUS Full-text
 DN 146:337575
 TI Arylacetic acids and related compounds as PPAR modulators and their preparation, pharmaceutical compositions and use in the treatment of PPAR-mediated diseases
 IN Lin, Jack; Womack, Patrick; Lee, Byunghun; Shi, Shenghua; Zhang, Chao; Artis, Dean R.; Ibrahim, Prabha N.; Wang, Weiru; Zuckerman, Rebecca
 PA Plexxikon, Inc., USA
 SO PCT Int. Appl., 239pp.
 CODEN: PIXKD2
 DT Patent
 LA English
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2007030567	A2	20070315	WO 2006-U934764	20060906
MO 2007030567	A3	20070621		

M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MM, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW
 RM: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,



L10 ANSWER 3 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
 AN 2007:88042 CAPLUS Full-text
 DN 146:172336
 TI Two photon-absorbing recording method, materials therefor, and recording/reading-out thereof
 IN Takizawa, Hiroo; Akiba, Masanaru
 PA Fuji Photo Film Co., Ltd., Japan
 SO Jpn. Kokai Tokkyo Koho, 103pp.
 CODEN: JKXXAF
 DT Patent
 LA Japanese
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI JP 2007017886	A	20070125	JP 2005-202022	20050711
PRAI JP 2005-202022		20050711		

OS MARPAT 146:172336

AB The materials contain (A) two photon-absorbing comps. and (B) components undergoing electron or energy transfer from excited state to exhibit (dis)coloration and are capable of recording through utilization of the change in refractive index, absorbance, or luminance, where A and/or B are oligomers or polymers. The compound A may be (merocyanine dyes, oxonol dyes, phthalocyanine dyes, azo dyes, and/or X102(CR104;CR103)m101CO(CR101;CR102)n101X101 [R101-R104 = H, substituent; n101, m101 = 0-4; n101 = m101 = 0; X101, X102 = aryl, heterocycle, Q (R105 = H, substituent; R106 = H, alk(en)yl, aryl, heterocycle; Z101 = 5- or 6-membered ring)]. Three-dimensional recording on the materials with laser light having longer wavelength than the linear absorption band of the compound A and with molar absorption coefficient ϵ 10, and their reading out by detecting reflectance or transmittance of reading light, are also claimed.


IT 520758-77-6

RI: TEM (Technical or engineered material use); USES (Uses)
 (recording components; three-dimensional optical recording materials containing oligomers or polymers with large two-photon absorption cross-sections)


RN 920758-77-6 CAPLUS

CN 2,4-Pentadienoic acid, 2-cyano-3-[3,5-dichloro-4-[(2-{4-[(diethylamino)sulfonyl]-2-nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-3-oxo-4-isoxazolyl)methoxy]phenyl]-, 6-[(2-methyl-1-oxo-2-propen-1-yl)oxy]hexyl ester, polymer with methyl 2-methyl-2-propenoate (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



CM 2
 CRN 80-62-6
 CMP C5 H8 O2




L10 ANSWER 4 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
 AN 2007:66228 CAPLUS [Full-text](#)
 DN 147:316984
 TI Docking and binding mode analysis of aryl diketoacids (ADK) at the active site of HCV RNA-dependent RNA polymerase
 AU Kim, J.; Chong, Y.
 CS Department of Biosciences and Biotechnology, Konkuk University, Seoul, 143-701, S. Korea

10581322-elected-species-final 7 of 76

were aligned by docking into the binding site, and a structure-based 3D-QSAR study was performed to correlate the biol. activities of ADKs with their three-dimensional structures. The CoMSIA model constructed by structure-based 3D-QSAR study could be successfully applied to predict the biol. activity of ADK analogs. The binding affinity of ADK analogs are found to be highly dependent upon the hydrogen bonding interaction as well as hydrophobic interaction around the aromatic ring of ADK analogs. In particular, the CoMSIA model proposes that the hydrophobic aromatic ring play a key role in determining the antiviral activity of ADK analogs. Thus, hydrophobic substituents around the aromatic ring reinforce hydrophobic interaction with the target enzyme, whereas the lack of aromatic substitution and thereby insufficient size of the inhibitor mol. can be primarily ascribed to their inability to bind to the hydrophobic binding site.

IT 932377-45-F
 RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)
 (structure-based 3D-QSAR(ComSIA) study on series of aryl diketoacids (ADK) as inhibitors of HCV RNA-dependent RNA polymerase)
 RN 932377-48-5 CAPLUS
 CN 2-Butenoic acid, 2-hydroxy-4-oxo-4-[3-[(5-phenyl-3-isoxazolyl)methoxy]phenyl]-, (2Z)- (CA INDEX NAME)

Double bond geometry as shown.




RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 6 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
 AN 2006:1278467 CAPLUS [Full-text](#)
 DN 146:27818
 TI Preparation of isoxazole derivatives as PPAR agonists
 IN Sugita, Kenichi; Kurose, Noriyuki; Katoaka, Mikayo; Setsukinal, Kenichi
 PA Shionogi and Co., Ltd., Japan
 SO Jpn. Kokai Tokkyo Koho, 123pp.
 CODEN: JKKXAP
 DT Patent
 LA Japanese
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2006328004	A	20061207	JP 2005-155739	20050527
JP 2005-155739		20050527		
MARPAT 146:27818				

AB The pharmacophore-guided docking study of aryl diketoacid (ADK) analogs revealed two distinctive hydrophobic binding sites (a pocket and a groove) around the UTP-binding site of hepatitis C virus (HCV) RNA-dependent RNA polymerase (RdRp). Interestingly, the hydrophobic binding sites have appropriate shape and size to specifically substituted aromatic rings, which suggests the specific role of substituents on the aromatic ring in determining the binding affinity of the ADK analog to the active site of the target enzyme. Binding mode anal. of ADK analogs with potent antiviral activity shows highly substituted aromatic rings map well onto the hydrophobic binding sites. For less active compds., their lack of aromatic substitution and thereby insufficient size can be primarily ascribed to their inability to bind to the hydrophobic binding site. The characteristic binding mode of ADK analogs proposed in this study provides a useful tool in designing a structure-activity relationship study of novel ADK analogs based on various aromatic substituents.


IT 268616-51-7P
 RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation) (docking and binding of aryl diketoacids (ADK) at the active site of HCV RNA-dependent RNA polymerase)
 RN 268616-51-7 CAPLUS
 CN 2-Butenoic acid, 2-hydroxy-4-oxo-4-[3-[(5-phenyl-3-isoxazolyl)methoxy]phenyl]- (CA INDEX NAME)



RE.CNT 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 5 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
 AN 2006:1335359 CAPLUS [Full-text](#)
 DN 146:374724
 TI A structure-based 3D-QSAR(ComSIA) study on a series of aryl diketoacids (ADK) as inhibitors of HCV RNA-dependent RNA polymerase
 AU Kim, Jinyoung; Han, Jin Hee; Chong, Youhoon
 CS Division of Biosciences and Biotechnology, Konkuk University, Seoul, 143-701, S. Korea
 SO Bulletin of the Korean Chemical Society (2006), 27(11), 1919-1922
 CODEN: BKCSDE; ISSN: 0253-2964
 PB Korean Chemical Society
 DT Journal
 LA English
 AB In this study, the hepatitis C virus (HCV) RNA-dependent RNA polymerase binding site used by the inhibitor aryl 4,γ-diketoacid (ADK) and its analogs was analyzed by using the crystal structure of rUTP-HCV RdRp complex (PDB ID 1GX6) and the structural similarity between rUTP and ADK. The ADK analogs


10581322-elected-species-final 8 of 76




AB Title compds. I [R1 = halo, OH, (un)substituted aryl, etc.; R2 - R10 = H, halo, (un)substituted alkyl, etc.; X1 = O, S, (un)substituted NH, etc.; X2 = (un)substituted CH2; X3 = carboxy, ester, etc.] and pharmaceutically acceptable salts and solvates thereof were prepared as peroxisome proliferator-activated receptor (PPAR) agonists. For instance, II was synthesized by thioetherification of the corresponding benzenethiol with 3-(chloromethyl)isoxazole, and ester hydrolysis of II led to the corresponding acid. Representative I showed PPAR agonistic activity with EC50 values of 1.0-28 nM. Therefore, the invented compds. are useful for the treatment of PPAR-related diseases.

IT 916240-92-1P 916240-94-3P 916240-95-5P
 J 916240-95-7P 916240-99-8P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (PPAR agonist; preparation of isoxazole derivs. as PPAR agonists)

RN 916240-92-1 CAPLUS
 CN Benzenepropanoic acid, 4-[[[4-(ethoxymethyl)-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-β-(trifluoromethyl)-, methyl ester (CA INDEX NAME)




RN 916240-94-3 CAPLUS
 CN Benzenepropanoic acid, β-(fluoromethyl)-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, methyl ester (CA INDEX NAME)



RN 916240-96-5 CAPLUS


CN Benzenepropanoic acid, beta-(difluoromethyl)-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, methyl ester (CA INDEX NAME)



RN 916240-98-7 CAPLUS


CN Benzenepropanoic acid, beta-methyl-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, methyl ester, (betaS)- (CA INDEX NAME)

Absolute stereochemistry.




RN 916240-99-8 CAPLUS

CN Benzenepropanoic acid, 4-[[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, methyl ester (CA INDEX NAME)




RN 916240-95-4 CAPLUS

CN Benzenepropanoic acid, beta-(difluoromethyl)-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, methyl ester (CA INDEX NAME)



RN 916240-97-6 CAPLUS


CN Benzenepropanoic acid, beta-(difluoromethyl)-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, methyl ester (CA INDEX NAME)



RN 916241-00-4 CAPLUS

CN Benzenepropanoic acid, 4-[[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-3-methoxy-beta-methyl-, (betaS)- (CA INDEX NAME)

(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-3-methoxy-beta-methyl-, methyl ester, (betaS)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry unknown.

IT 916085-42-2 CP 916240-92-2P 916240-55-4P

916240-57-6P 916241-00-4E 916241-01-5P

916241-02-6P 916241-03-7P 916241-04-9P

916241-05-9P 916241-06-0P 916241-07-1P

916241-08-4P 916241-09-3P 916241-10-5P


916241-11-7P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

RN 916085-42-2 CAPLUS


CN Benzenepropanoic acid, beta-methyl-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, (betaS)- (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 916240-93-2 CAPLUS

CN Benzenepropanoic acid, 4-[[[4-(ethoxymethyl)-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-beta-(trifluoromethyl)- (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry unknown.


RN 916241-01-5 CAPLUS

CN Benzenepropanoic acid, 4-[[[5-(2,6-difluorophenyl)-4-methyl-3-isoxazolyl]methyl]thio]-beta-(trifluoromethyl)- (CA INDEX NAME)




RN 916241-02-6 CAPLUS

CN Benzenepropanoic acid, 4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-beta-(trifluoromethyl)- (CA INDEX NAME)




RN 916241-03-7 CAPLUS


CN Benzenepropanoic acid, 4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-beta-(trifluoromethyl)- (CA INDEX NAME)




RN 916241-04-8 CAPLUS
CN Benzenepropanoic acid, 4-[[[4-[(ethoxymino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-β-(trifluoromethyl)- (CA INDEX NAME)



RN 916241-05-9 CAPLUS
CN Benzenepropanoic acid, 4-[[[4-(ethoxymethyl)-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-β-(trifluoromethyl)- (CA INDEX NAME)




RN 916241-06-0 CAPLUS
CN Benzenepropanoic acid, 4-[[[5-(4-chlorophenyl)-4-methyl-3-isoxazolyl]methyl]thio]-β-(trifluoromethyl)- (CA INDEX NAME)




RN 916241-10-6 CAPLUS
CN Benzenepropanoic acid, 3-methoxy-β-methyl-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-, (βS)- (CA INDEX NAME)

Absolute stereochemistry.




RN 916241-11-7 CAPLUS
CN Benzenepropanoic acid, 4-[[[4-(ethoxymethyl)-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-3-methoxy-β-methyl-, (βS)- (CA INDEX NAME)


Absolute stereochemistry.




L10 ANSWER 7 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
AN 2006125037 CAPLUS Full-text
DN 146:33027
TI Pharmaceutical composition comprising vitamin K



RN 916241-07-1 CAPLUS
CN Benzenepropanoic acid, 3-methoxy-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-β-(trifluoromethyl)- (CA INDEX NAME)



RN 916241-08-2 CAPLUS
CN Benzenepropanoic acid, 4-[[[4-[(ethoxymino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-β-(trifluoromethyl)- (CA INDEX NAME)




RN 916241-09-3 CAPLUS
CN Benzenepropanoic acid, 4-[[[4-(ethoxymethyl)-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-β-(trifluoromethyl)- (CA INDEX NAME)

IN Inoue, Satoshi; Sato, Seiji; Kyokawa, Yoshimasa; Sugita, Ken-Ichi; Torii, Mikinori
PA Shionogi & Co., Ltd., Japan
SO PCT Int. Appl., 91pp.
CODEN: PIXXD2
DT Patent
LA Japanese
FAN.CWT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2006126541	A1	20061130	WO 2006-JP10249	20060523
M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MM, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MM, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				

PRA1 JP 2005-155837 A 20050527
AB It is found that a compound having a PPAR δ agonistic activity induces abnormal blood coagulation or a muscular disorder. A pharmaceutical composition comprising the combination of a compound having a PPAR δ agonistic activity and a vitamin K can prevent the abnormal blood coagulation. A pharmaceutical composition comprising a vitamin K can prevent the muscular disorder.
IT 254013-20-0 254014-61-2 915728-67-9
915730-23-1 915739-22-0 916085-42-2
916085-13-3 916085-10-8
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(pharmaceutical composition comprising vitamin K)


RN 854013-20-0 CAPLUS
CN 2-Propenoic acid, 2-chloro-3-[3-methoxy-4-[[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]phenyl]- (CA INDEX NAME)




RN 854014-61-2 CAPLUS
CN Benzenepropanoic acid, 4-[[[4-[(ethoxymino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-3-methoxy-β-methyl- (CA INDEX NAME)





RN 915788-50-0 CAPLUS
CN Benzeneacetic acid, 3-[[4-methyl-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-51-1 CAPLUS
CN Benzeneacetic acid, 3-[[4-(ethoxymethyl)-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-52-2 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(cyclopropylmethoxy)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)



RN 915788-57-7 CAPLUS
CN Benzeneacetic acid, 4-chloro-3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-58-8 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-4-methoxy- (CA INDEX NAME)




RN 915788-59-9 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-5-methoxy- (CA INDEX NAME)


RN 915788-53-3 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-54-4 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]thio- (CA INDEX NAME)




RN 915788-55-5 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-4-fluoro- (CA INDEX NAME)




RN 915788-56-6 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]thio-4-fluoro- (CA INDEX NAME)




RN 915788-60-2 CAPLUS
CN Benzeneacetic acid, 2-chloro-5-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)



RN 915788-61-3 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-2-methyl- (CA INDEX NAME)




RN 915788-63-5 CAPLUS
CN Benzeneacetic acid, 3-chloro-5-[[4-[(ethoxymethyl)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]thio- (CA INDEX NAME)




RN 915788-64-6 CAPLUS

CN Benzeneacetic acid, 3-[[[5-(4-chlorophenyl)-4-[[[2-(2-fluoroethoxy)imino]methyl]-3-isoxazolyl]methyl]thio]- (CA INDEX NAME)




RN 915788-65-7 CAPLUS

CN Benzeneacetic acid, 3-[[[5-(4-chlorophenyl)-4-[[[2-(2-fluoroethoxy)imino]methyl]-3-isoxazolyl]methoxy]-4-methyl- (CA INDEX NAME)




RN 915788-66-8 CAPLUS

CN Benzeneacetic acid, 3-[[[5-(4-chlorophenyl)-4-[[[2-(2-fluoroethoxy)imino]methyl]-3-isoxazolyl]methyl]thio]-4-methyl- (CA INDEX NAME)




RN 915788-70-4 CAPLUS

CN Benzeneacetic acid, 3-chloro-5-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-72-6 CAPLUS

CN Benzeneacetic acid, 3-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-73-7 CAPLUS

CN Benzeneacetic acid, 4-chloro-5-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-67-9 CAPLUS

CN Benzeneacetic acid, 4-chloro-3-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-68-0 CAPLUS

CN Benzeneacetic acid, 4-chloro-3-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-69-1 CAPLUS

CN Benzeneacetic acid, 3-methyl-5-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-74-8 CAPLUS

CN Benzeneacetic acid, 3-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-75-9 CAPLUS

CN Benzeneacetic acid, 3-chloro-5-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-76-0 CAPLUS

CN Benzeneacetic acid, 3-chloro-5-[[[4-[[[2-(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915789-77-1 CAPLUS

CN Benzenecetic acid, 3-[[4-[(cyclopropylmethoxy)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-u,u-difluoro- (CA INDEX NAME)




RN 915788-78-2 CAPLUS

CN Benzenecetic acid, 3-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]-u,u-difluoro- (CA INDEX NAME)




RN 915788-79-3 CAPLUS

CN Benzenecetic acid, 4-chloro-3-[[4-[(cyclobutyloxy)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-83-9 CAPLUS

CN Benzenecetic acid, 3-[[4-[(cyclobutyloxy)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methyl]thio]-4-methyl- (CA INDEX NAME)




RN 915788-84-0 CAPLUS

CN Benzenecetic acid, 3-[[4-[(cyclobutyloxy)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]-4-methyl- (CA INDEX NAME)




RN 915788-86-2 CAPLUS

CN Benzenecetic acid, 3-[[4-[(2-chloroethoxy)imino]methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]- (CA INDEX NAME)




RN 915788-80-6 CAPLUS

CN Benzenecetic acid, 3-[[4-[(cyclobutyloxy)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methyl]thio]- (CA INDEX NAME)




RN 915788-81-7 CAPLUS

CN Benzenecetic acid, 4-chloro-3-[[4-[(cyclobutyloxy)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methyl]thio]- (CA INDEX NAME)




RN 915788-82-8 CAPLUS

CN Benzenecetic acid, 3-[[4-[(cyclobutyloxy)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-87-3 CAPLUS

CN Benzenecetic acid, 3-[[4-[[2-[(1,1-dioxido-4-thiomorpholinyl)ethoxy]imino]methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio]- (CA INDEX NAME)




RN 915788-88-4 CAPLUS

CN Benzenecetic acid, 3-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-4-methyl- (CA INDEX NAME)




RN 915788-89-5 CAPLUS


CN Benzenecetic acid, 3-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-5-fluoro- (CA INDEX NAME)




RN 915788-90-8 CAPLUS
CN Benzeneacetic acid, 5-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-2-fluoro- (CA INDEX NAME)




RN 915788-91-9 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-2-fluoro- (CA INDEX NAME)




RN 915788-92-0 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-2-methoxy- (CA INDEX NAME)




RN 915788-96-4 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-2-thio- (CA INDEX NAME)




RN 915788-97-5 CAPLUS
CN Benzeneacetic acid, 3-chloro-5-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-2-thio- (CA INDEX NAME)




RN 915788-98-6 CAPLUS
CN Benzeneacetic acid, 4-methyl-3-[[4-[(propoxyimino)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methoxy]- (CA INDEX NAME)




RN 915788-93-1 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(cyclopropylmethoxy)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio- (CA INDEX NAME)




RN 915788-94-2 CAPLUS
CN Benzeneacetic acid, 3-[[4-[[[(1-methylethylidene)amino]oxy]methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio- (CA INDEX NAME)




RN 915788-95-3 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxymethyl)-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methyl]thio- (CA INDEX NAME)




RN 915788-99-7 CAPLUS
CN Benzeneacetic acid, 3-chloro-5-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methyl]thio- (CA INDEX NAME)



RN 915789-00-3 CAPLUS
CN Benzeneacetic acid, 3-chloro-5-[[4-[[[(2-fluoroethoxy)imino]methyl]-5-[4-(trifluoromethoxy)phenyl]-3-isoxazolyl]methyl]thio]- (CA INDEX NAME)




IT 915789-10-5 915789-11 4P 915789 12 7P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of arylacetate derivs. containing isoxazole moiety as PPAR agonists)
RN 915789-10-5 CAPLUS
CN Benzeneacetic acid, 3-[[4-[(ethoxyimino)methyl]-5-[4-(trifluoromethyl)phenyl]-3-isoxazolyl]methoxy]-5-methyl-, methyl ester (CA INDEX NAME)




RN 915789-11-6 CAPLUS

CN Benzeneacetic acid, 3-[[[4-formyl-5-{4-(trifluoromethyl)phenyl}-3-isoxazolyl]methyl]thio]-, methyl ester (CA INDEX NAME)



RN 915789-12-7 CAPLUS

CN Benzeneacetic acid, 3-[[[4-[[[2-fluoromethoxy]imino]methyl]-5-{4-(trifluoromethyl)phenyl}-3-isoxazolyl]methyl]thio]-, methyl ester (CA INDEX NAME)

RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 9 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN

AN 2006:1001108 CAPLUS [Full-text](#)
DN 146:208TI 3,4,5-Trisubstituted isoxazoles as novel PPAR δ agonists. Part 2AU Epple, Robert; Azimioara, Mihai; Russo, Ross; Xie, Yongping; Wang, Xing; Cow, Christopher; Wityak, John; Karanewsky, Don; Bursulaya, Badry; Kreusch, Andreas; Tuntland, Tove; Gerken, Andrea; Iskandar, Maya; Saez, Enrique; Martin Seidel, H.; Tian, Shin-Shay
CS Department of Medicinal Chemistry, The Genomics Institute of the Novartis Research Foundation, San Diego, CA, 92121, USASO Bioorganic & Medicinal Chemistry Letters (2006), 16(21), 5488-5492
CODEN: BMCL68; ISSN: 0960-894X


PB Elsevier Ltd.

DT Journal

LA English

OS CASREACT 146:208

GI

AB A series of PPAR δ -selective agonists was investigated and optimized for a favorable in vivo pharmacokinetic profile. Isoxazole LC1765 (I) was a potent and selective PPAR δ agonist with good in vivo PK properties in mouse (Cmax = 5.1 μ M, t1/2 = 3.1 h). LC1765 regulated expression of genes involved in energy homeostasis in relevant tissues when dosed orally in C57BL6 mice. A co-crystal structure of compound LC1765 and the LBD of PPAR δ is discussed.


IT 21:194-71:7P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(3,4,5-trisubstituted isoxazoles as novel PPAR δ agonists and structure activity relations)

RN 915194-71-7 CAPLUS

CN Benzeneacetic acid, 3-chloro-4-[[[4-[[[2-(2,4-dichlorophenoxy)ethyl]amino]carbonyl]-5-phenyl]-3-isoxazolyl]methoxy]-, methyl ester (CA INDEX NAME)




IT 21:194-71:7P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(3,4,5-trisubstituted isoxazoles as novel PPAR δ agonists and structure activity relations)


RN 927178-35-6 CAPLUS

CN 4-isoxazolecarboxylic acid, 3-[[[2-chloro-4-(2-methoxy-2-oxoethyl)phenoxy]methyl]-5-phenyl]-, 1,1-dimethylethyl ester (CA INDEX NAME)



RN 927178-36-7 CAPLUS


CN Benzeneacetic acid, 3-chloro-4-[[[4-(chlorocarbonyl)-5-phenyl]-3-isoxazolyl]methoxy]-, methyl ester (CA INDEX NAME)



RN 927178-54-9 CAPLUS

CN Benzeneacetic acid, 3-chloro-4-[[[4-[[[2-(2,4-dichlorophenoxy)ethyl]amino]carbonyl]-5-phenyl]-3-isoxazolyl]methoxy]-, methyl ester (CA INDEX NAME)

PAGE 1-A




PAGE 2-A

Cl

RN 927192-17-4 CAPLUS

CN 4-isoxazolecarboxylic acid, 3-[[[2-chloro-4-(2-methoxy-2-oxoethyl)phenoxy]methyl]-5-phenyl]- (CA INDEX NAME)

RE.CNT 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 10 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN

AN 2006:914124 CAPLUS [Full-text](#)
DN 145:325062

TI Holographic optical information recording medium with high sensitivity having cholesteric liquid crystal layer
 IN Takizawa, Hiroo
 PA Fuji Photo Film Co., Ltd., Japan
 SO Jpn. Kokai Tokkyo Koho, 67pp.
 CODEN: JKXXAF
 DT Patent
 LA Japanese
 FAN. CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2006235385	A	20060907	JP 2005-51772	20050225
PRAI JP 2005-51772		20050225		

AB Disclosed is a holog. optical information recording medium comprising (a) a translucent substrate, (b) a holog. recording layer formed on the substrate, and (c) a filter layer interposed between (a) and (b) capable of transmitting a 1st wavelength and reflecting a 2nd wavelength, wherein said recording layer includes an optical refractive index modulation component capable of recording interference fringes as a refractive index modulation by various reaction means.


IT 507159-03-5
 RL: DEV (Device component use); USES (Uses)
 (Holog. optical information recording disk with high sensitivity having cholesteric liquid crystal layer)

RN 907199-03-5 CAPLUS

CN 2-Propenoic acid, 2-cyano-3-[3,5-dichloro-4-[[2-[4-[(diethylamino)sulfonyl]-2-nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-3-oxo-4-isoxazolyl]methoxy]phenyl]-, ethyl ester (CA INDEX NAME)



PAGE 1-A



PAGE 2-A

L10 ANSWER 12 OF 25 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2006:913844 CAPLUS [Full-text](#)
 DN 145:325042
 TI Hologram information recording method for high sensitivity and high density

IN Takizawa, Hiroo
 PA Fuji Photo Film Co., Ltd., Japan
 SO Jpn. Kokai Tokkyo Koho, 71pp.
 CODEN: JKXXAF

DT Patent
 LA Japanese
 FAN. CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2006235209	A	20060907	JP 2005-49176	20050224
PRAI JP 2005-49176		20050224		

AB A hologram information recording method utilizes an optical refractive index-modulating of an information recording layer by (1) a color development reaction, (2) a color development reaction amplified by a self-sensitization reaction with a coloring material of a latent image, (3) a polymerization reaction sensitized by a coloring material of a latent image, (4) an alignment change in a compound having a birefringence, (5) a dye discoloration reaction, or (6) a latent image-sensitized polymerization reaction sensitized by a latent image of a residual of a discolorable dye.

IT 907199-03-5
 RL: DEV (Device component use); USES (Uses)
 (holog. recording material; hologram information recording method for high sensitivity and high d.)

RN 907199-03-5 CAPLUS

CN 2-Propenoic acid, 2-cyano-3-[3,5-dichloro-4-[[2-[4-[(diethylamino)sulfonyl]-2-nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-3-oxo-4-isoxazolyl]methoxy]phenyl]-, ethyl ester (CA INDEX NAME)



PAGE 2-A

L10 ANSWER 11 OF 25 CAPLUS COPYRIGHT 2008 ACS on STN

AN 2006:913846 CAPLUS [Full-text](#)

DN 145:325043

TI Hologram information recording method for high sensitivity and high density

IN Takizawa, Hiroo
 PA Fuji Photo Film Co., Ltd., Japan
 SO Jpn. Kokai Tokkyo Koho, 72pp.
 CODEN: JKXXAF

DT Patent
 LA Japanese
 FAN. CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2006235210	A	20060907	JP 2005-49177	20050224
PRAI JP 2005-49177		20050224		

AB A hologram information recording method utilizes an optical refractive index-modulating of an information recording layer by (1) a color development reaction, (2) a color development reaction amplified by a self-sensitization reaction with a coloring material of a latent image, (3) a polymerization reaction sensitized by a coloring material of a latent image, (4) an alignment change in a compound having a birefringence, (5) a dye discoloration reaction, or (6) a latent image-sensitized polymerization reaction sensitized by a latent image of a residual of a discolorable dye.


IT 507199-03-5
 RL: DEV (Device component use); USES (Uses)
 (holog. recording material; hologram information recording method for high sensitivity and high d.)

RN 907199-03-5 CAPLUS

CN 2-Propenoic acid, 2-cyano-3-[3,5-dichloro-4-[[2-[4-[(diethylamino)sulfonyl]-2-nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-3-oxo-4-isoxazolyl]methoxy]phenyl]-, ethyl ester (CA INDEX NAME)



PAGE 1-A



PAGE 2-A

L10 ANSWER 13 OF 25 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2006:890375 CAPLUS [Full-text](#)
 DN 145:281119
 TI Method and apparatus for formation of hologram for readout of three-dimensional images


IN Takizawa, Hiroo
 PA Fuji Photo Film Co., Ltd., Japan
 SO Jpn. Kokai Tokkyo Koho, 80pp.
 CODEN: JKXXAF

DT Patent
 LA Japanese
 FAN. CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2006227067	A	20060831	JP 2005-37603	20050215
PRAI JP 2005-37603		20050215		

AB The apparatus for recording interference fringes on recording media for generation of reconstructed light according to desirable three-dimensional images when irradiating reference light, consists of a head for irradiation of several light fluxes for recording interference fringes, and a means for changing relative positions between the head and the recording media, wherein the interference fringes are recorded as refractive index modulation by polymerization, color development, self-sensitization and amplification color development from latent images, etc. The recording media show high diffraction efficiency and low shrinkage, and are useful for multiple recording.

IT 2006:851010 CAPLUS [Full-text](#)
 RL: TEM (Technical or engineered material use); USES (Uses)
 (interference fringe-recording component; method and apparatus for
 formation of hologram for readout of three-dimensional images)
 RN 907199-03-5 CAPLUS
 CN 2-Propenoic acid, 2-cyano-3-[(3,5-dichloro-4-[[2-[[4-
 [(diethylamino)sulfonyl]-2-nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-
 3-oxo-4-isoxazoly]methoxy]phenyl]-, ethyl ester (CA INDEX NAME)




PAGE 1-A




PAGE 2-A

RN 907199-11-5 CAPLUS
 CN 2-Propenoic acid, 2-cyano-3-[(4-[[2-[[4-[(diethylamino)sulfonyl]-2-
 nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-3-oxo-4-
 isoxazoly]methoxy]phenyl]-, ethyl ester (CA INDEX NAME)




PAGE 1-A



PAGE 2-A

RN 907199-12-6 CAPLUS
 CN 2-Propenoic acid, 2-cyano-3-[(4-[[2-[[4-[(diethylamino)sulfonyl]-2-
 nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-3-oxo-4-isoxazoly]methoxy]-
 3,5-dimethylphenyl]-, ethyl ester (CA INDEX NAME)



PAGE 1-A


L10 ANSWER 14 OF 25 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2006:851010 CAPLUS [Full-text](#)
 DN 145:281118
 TI Hologram recording material, hologram recording method, and optical
 recording medium
 IN Takizawa, Hiroo
 PA Fujii Photo Film Co., Ltd., Japan
 SO U.S. Pat. Appl. Publ., 69pp.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN, CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI US 2006189790	A1	20060824	US 2006-359566	20060223
JP 2006235087	A	20060907	JP 2005-47609	20050223
PRAI JP 2005-47609	A	20050223		


AB A hologram recording material is provided and has: a sensitizing dye absorbing light upon hologram exposure to generate an excited state thereof, and an interference fringes-recording component capable of causing color development reaction or discoloration by an electron or energy transfer (movement) form the excited state to record interference fringes providing a refractive index modulation. The sensitizing dye or the interference fringes-recording component is a polymer or an oligomer. The recording material is applied to high d. optical recording medium, three-dimensional display, holog. optical element, etc.

IT 2006:851010 CAPLUS [Full-text](#)
 RL: TEM (Technical or engineered material use); USES (Uses)
 (interference fringes-recording component in hologram recording
 material)
 RN 906543-53-1 CAPLUS
 CN 2-Propenoic acid, 2-methyl-, methyl ester, polymer with
 6-[(2-methyl-1-oxo-2-propenyl)oxy]hexyl 2-cyano-3-[(3,5-dichloro-4-[[2-[[4-
 [(diethylamino)sulfonyl]-2-nitrophenyl]-5-(1,1-dimethylethyl)-2,3-dihydro-
 3-oxo-4-isoxazoly]methoxy]phenyl]-2-propenoate (9C1) (CA INDEX NAME)

CM 1
 CRN 906543-52-0
 CMF C38 H44 Cl2 N4 O11 S




PAGE 1-A



PAGE 2-A

CM 2
 CRN 80-62-6
 CMF C5 H8 O2



L10 ANSWER 15 OF 25 CAPLUS COPYRIGHT 2008 ACS on STN
 AN 2006:364586 CAPLUS [Full-text](#)
 DN 144:412487
 TI Isoxazole and isothiazole compounds as PPAR α agonists, their
 preparation, pharmaceutical compositions, and use in therapy
 IN Madhavan, Gurrām Range, Iqbal, Javed, Bhuniya, Debnath, Das, Seibal Kumar,
 Sharma, Sudhir Kumar, Chakrabarti, Ranjan
 PA Dr. Reddy's Laboratories Ltd., India; Dr. Reddy's Laboratories, Inc.
 SO PCT Int. Appl., 65 pp.
 CODEN: PIXXD2
 DT Patent
 LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006042245	A1	20060420	WO 2005-US36474	20051011




W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

PRAI IN 2004-CH1051 A 20041011

OS CASREACT.144:412487; MARPAT 144:412487

GI



AB The invention relates to isoxazoles and related compds. of formula I, which are peroxisome proliferator-activated receptor (PPAR) agonists, specifically the PPAR α subtype. In compds. I, Ar1 is (un)substituted aryl or (un)substituted heteroaryl; Ar2 is (un)substituted aryl; W is O, S, or CH₂; X is O or S; Y is O, S, CH₂, or NR₅, where R₅ is H, alkyl, or cycloalkyl; n is 0 or 1; m is 0-6; R1 and R2 are independently selected from H, OH, halo, (un)substituted alkyl, (un)substituted cycloalkyl, (un)substituted aryl, (un)substituted heteroaryl, (un)substituted aryloxy, (un)substituted heteroaryl, (un)substituted heterocyclyl, (un)substituted heteroalkyl, or R1 and R2 together form an (un)substituted 5- or 6-membered ring, optionally containing one or two heteroatoms selected from O, S, and N; and R3 and R4 are independently selected from H, (un)substituted alkyl, (un)substituted cycloalkyl, (un)substituted aryl, (un)substituted heteroaryl, (un)substituted heterocyclyl, and (un)substituted heteroalkyl. The invention also relates to the preparation of I, pharmaceutical compds. comprising a compound of formula I and one or more pharmaceutically acceptable

excipients, as well as to the use of the compns. for the treatment of diseases or disorders that respond to PPAR α activation. Cyclization of N-hydroxybenzenecarboximidoyl chloride with 1-methoxy-4-(pent-4-ynyl)benzene and demethylation gave isoxazole II, which underwent alkylation with Et 2-bromoisobutyrate and ester hydrolysis to give isoxazole III. The compds. of the invention act as agonists of PPAR α , e.g., compound III expresses 1.5-fold, 4.8-fold, and 5.9-fold activation of luciferase (mediated by PPAR α) compared with untreated cells at concns. of 1 μ M, 10 μ M, and 50 μ M, resp.


IT 883750-38-7P 883750-39-8P

RI: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of isoxazoles and isothiazoles as PPAR α agonists)


RN 883750-38-7 CAPLUS

CN Benzenepropanoic acid, α ,2-dimethyl-4-[(3-phenyl-5-isoxazolyl)methoxy]- (CA INDEX NAME)



RN 883750-39-8 CAPLUS

CN Benzenepropanoic acid, α -methyl- α -phenoxy-4-[(3-phenyl-5-isoxazolyl)methoxy]- (CA INDEX NAME)



RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 16 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN

AN 2005:1026833 CAPLUS Full-text

DN 143:326090

TI Preparation of arylmethoxyphenyl-alkylcarboxylic acids and related derivatives for use in treating metabolic disorders

IN Akerman, Michelle; Houze, Jonathan; Lin, Daniel C. H.; Liu, Jiwen; Luo, Jian; Medina, Julio C.; Qiu, Mei; Reagan, Jeffrey D.; Sharma, Rajiv; Shuttleworth, Stephen J.; Sun, Ying; Zhang, Jian; Zhu, Liusheng

PA Angen Inc., USA; et al.

SO PCT Int. Appl., 163 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005086661	A2	20050922	WO 2005-US5815	20050224
WO 2005086661	A3	20060504		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BF, BJ, CF, CG, CI, CM, CN, CO, CR, CU, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

AU 2005220728 A2 20050922 AU 2005-220728 20050224

AU 2005220728 A1 20050922

CA 2558585 A1 20050922 CA 2005-2558585 20050224

EP 1737809 A2 20070103 EP 2005-723623 20050224

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, LV, MK, YU

CN 1946666 A 20070411 CN 2005-80012709 20050224

BR 2005008098 A 20070717 BR 2005-8098 20050224

JP 2007525516 T 20070906 JP 2007-500959 20050224

US 2006004012 A 20060105 US 2005-67377 20050225

MX 2006PA09793 A 20061030 MX 2006-PA9793 20060828

US 2007142384 A1 20070621 US 2006-591214 20060828

KR 2007004769 A 20070109 KR 2006-719713 20060922

IN 2006DN05525 A 20070817 IN 2006-DN5525 20060922

NO 200604362 A 20061122 NO 2006-4362 20060926


PRAI US 2004-548741P P 20040227

US 2004-601579P P 20040812

WO 2005-US5815 W 20050224

OS MARPAT 143:326090

GI



AB Title compds. O-L1-P-L2-M-X-L3-A [O = H, (hetero)aryl, alkyl, etc.; L1 = bond, alkylene, heteroalkylene, O, etc.; P = (hetero)aromatic, cycloalkylene, etc.; L2 = bond, alkylene, heteroalkylene, etc.; M = (hetero)aromatic, cycloalkylene, arylalkylene, etc.; X = divalent alkyl, (un)substituted-N; O, SOO-2; L3 = bond, alkylene, heteroalkylene, etc.; A = COOH, tetrazolyl, SO₃H, PO₃H₂, etc.; 1] are prepared For instance, (S)-3-[4-((4'-trifluoromethyl-1,1'-biphenyl-3-yl)methoxy)phenyl]hexan-4-ynoic acid (II) is prepared in 5 steps from (S)-3-(4-hydroxyphenyl)hexan-4-ynoic acid Me ester (preparation given), 4-(trifluoromethyl)phenylboronic acid and 3-bromobenzoic acid. II has an EC₅₀ < 0.1 μ M for human G protein-coupled receptor GPR40. I are useful for the treatment of type II diabetes.


IT 855232-99-1P

RI: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of arylmethoxyphenyl-alkylcarboxylic acids and related derivs. as GPCR40 ligands for use in treating metabolic disorders)

RN 855232-99-1 CAPLUS

CN Benzenepropanoic acid, 4-[(3-phenyl-5-isoxazolyl)methoxy]- β -1-propynyl- (9CI) (CA INDEX NAME)



L10 ANSWER 17 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN

AN 2005:980049 CAPLUS Full-text

DN 143:431884

TI Simple but Highly Effective Three-Dimensional Chemical-Feature-Based Pharmacophore Model for Diketo Acid Derivatives as Hepatitis C Virus RNA-Dependent RNA Polymerase Inhibitors

AU Di Santo, Roberto; Permeleglia, Maurizio; Perrone, Marco; Paneni, Maria Silvia; Costi, Roberta; Artico, Marino; Roux, Alessandra; Gabriele, Mirko; Tardif, Keith D.; Siddiqui, Aileen; Prisci, Sabrina

CS Istituto Pasteur-Fondazione Cenci Bolognietti-Dipartimento di Studi Farmaceutici, University of Rome La Sapienza, Rome, I-00185, Italy

SO Journal of Medicinal Chemistry (2005), 48(20), 6304-6314
CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society


DT Journal

LA English

AB A mol. modeling strategy using aryl diketo acid (ADK) derivs. recently reported in the literature as hepatitis C virus (HCV) polymerase inhibitors was designed. A 3D chemical-feature-based pharmacophore model was developed using Catalyst software, which produced 10 pharmacophore hypotheses. The top-ranked one (Hypo 1), characterized by a high correlation coefficient ($r = 0.965$), consisted of two hydrogen bond acceptors, one neg. ionizable moiety, and two hydrophobic aroms. This model was used to predict the anti-RNA-dependent RNA polymerase (anti-RdRp) activity of 6-(1-arylmethylpyrrol-2-yl)-1,4-dioxo-5-hexenoic acids and other ADK derivs. previously synthesized in our labs. as HIV-1 integrase inhibitors. Furthermore, the expl. IC50 values of 9 compds., tested in vitro against recombinant HCV polymerase, were compared with the corresponding values predicted using Hypo1. A good agreement between expl. and simulated data was obtained. The results demonstrate that the hypothesis derived in this study can be considered to be a useful tool in designing new leads based on ADK scaffolds as HCV RdRp inhibitors.

IT **REMARKS:**
 RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (pharmacophore model for diketo acid derivs. as hepatitis C virus RNA-dependent RNA polymerase inhibitors)

RN 858616-51-7 CAPLUS
 CN 2-Butenoic acid, 2-hydroxy-4-oxo-4-[3-[(5-phenyl-3-isoxazolyl)methoxy]phenyl]- (CA INDEX NAME)



RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 18 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN

AN 2005:158622 CAPLUS Full-text

DN 142:279952

TI Preparation of aralkanoates as inhibitors of prostaglandin and leukotriene production.

IN Shoda, Motoshi; Kuriyama, Hiroshi

PA Asahi Kasei Pharma Corporation, Japan

SO PCT Int. Appl., 687 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 4

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
MO 2005016862	A1	20050224	WO 2004-JP11952	20040813
M:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RM:	BW, GH, GM, KE, LS, MM, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2004265191 A1 20050224 AU 2004-265191 20040813
 CA 2535665 A1 20050224 CA 2004-2535665 20040813
 WO 2005016862 A1 20050224 WO 2004-XA11952 20040813

M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RM: BW, GH, GM, KE, LS, MM, MZ, NA, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

MO 2005016862 A1 20050224 WO 2004-XB11952 20040813

M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RM: BW, GH, GM, KE, LS, MM, MZ, NA, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

MO 2005016862 A1 20050224 WO 2004-XC11952 20040813

M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, ME, MK, MN, MW, MX, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RM: BW, GH, GM, KE, LS, MM, MZ, NA, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

EP 1660427 A1 20060531 EP 2004-771913 20040813

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK

CN 101031539 A 20070905 CN 2004-80024789 20040813

JP 2007280362 T 20071011 JP 2006-519267 20040813


MX 2006PA01739 A 20060512 MX 2006-PA1739 20060214

US 2007213333 A1 20070913 US 2007-568185 20070122

PRAI JP 2003-293590 A 20030814
 US 2003-495734P P 20030818
 WO 2004-JP11952 W 20040813



OS CASREACT 142:279952; MARPAT 142:279952

GI



IT 7389913-52-42, and 7389913-52-43 (no data).
7389913-52-47, 3-[2-Methyl-4-[1-(4-methyl-1-(4-
trifluoromethylphenyl)isoxazol-5-yl)ethoxy]phenyl]propionic acid
7389913-52-52, 3-[2-Methyl-4-[4-methyl-1-(4-
trifluoromethylphenyl)isoxazol-5-ylmethoxy]phenyl]propionic acid
RL: PAC (Pharmacological activity), SPN (Synthetic preparation), THU
(Therapeutic use), BIOL (Biological study), PREP (Preparation), USES
(Uses)
[PPAR modulators, preparation of pyrazoles and analogs as PPAR modulators]

RN 728913-82-4 CAPLUS
CN Benzenepropanoic acid, 2-methyl-4-[1-[4-methyl-3-[4-(trifluoromethyl)phenyl]-5-isoxazolyl]ethoxy]- (CA INDEX NAME)




and UMR, details of the preparation. The molecular weight is also given.


IT 723332-44-9P 623332-45-7P 623332-46-8P 623332-47-9P 623332-48-2P 623332-49-2P 623332-44-9P 623332-6A-1P 623332-71-0P 623332-75-2P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses).


RN 628332-46-7 CAPLUS
CN Benzeneacetic acid, 3-[3-[3-[4-(trifluoromethyl)phenyl]-5-isoxazolyl]propoxy]- (CA INDEX NAME)



RN 628332-48-9 CAPLUS
CN Benzenepropanoic acid, 4-[3-[3-[4-(trifluoromethyl)phenyl]-5-
isoxazolyl]propoxy]- (CA INDEX NAME)




RN 628332-57-0 CAPLUS
CN Benzenecetic acid, 4-[4-[3-[4-(trifluoromethyl)phenyl]-5-
isoxazolyl]butoxy]- (CA INDEX NAME)




RN 628332-59-2 CAPLUS

CN Benzenepropanoic acid, 4-[4-[3-[4-(trifluoromethyl)phenyl]-5-isoxazolyl]butoxy]- (CA INDEX NAME)



RN 628332-64-9 CAPLUS

CN Benzenepropanoic acid, 3-[4-[3-[4-(trifluoromethyl)phenyl]-5-isoxazolyl]butoxy]- (CA INDEX NAME)




RN 628332-66-1 CAPLUS

CN Benzenepropanoic acid, 2-[4-[3-[4-(trifluoromethyl)phenyl]-5-isoxazolyl]butoxy]- (CA INDEX NAME)


AN 2003:154240 CAPLUS Full-text
 DN 138:198669
 TI FXR NR1H4 nuclear receptor binding compounds
 IN Bauer, Ulrike; Cheruvallath, Zach; Deuschle, Ulrich; Dneprovskaja, Elena; Gahman, Tim; Giegrich, Kristina; Hanecak, Ronnie; Hebert, Normand; Kiely, John; Kober, Ingo; Kogel, Manfred; Kranz, Harald; Kremoser, Claus; Lee, Matthew; Olte, Kerstin; Sage, Carlton; Sud, Manish
 PA Lion Bioscience AG, Germany
 SO PCT Int. Appl., 53 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN CNT 5

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003015771	A1	20030227	WO 2002-US25437	20020813
W: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MM, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RM: GH, GM, KE, LS, MN, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1285914	A1	20030226	EP 2001-119473	20010813
EP 1285914	B1	20071219		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
US 2003187042	A1	20031002	US 2002-185721	20020701
US 7034046	B2	20060425		
AU 2002319805	A1	20030303	AU 2002-319805	20020813
EP 1423111	A1	20040602	EP 2002-750473	20020813
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
PRAI EP 2001-119473	A	20010813		
US 2002-185721	A	20020701		
WO 2002-US25437	W	20020813		
OS MARPAT 138:198669				
AB The present invention relates to compds. according to the general formula (I) which bind to the nuclear receptor, NR1H4 (farnesoid X receptor α), and act as agonists, antagonists or mixed agonists/antagonists of the NR1H4 receptor. The invention further relates to the treatment of diseases and/or conditions through binding of the nuclear receptor by the compds. It was further an object of the invention to provide for compds. which may be used for the manufacture of a medicament for the treatment of cholesterol or bile acid associated conditions or diseases. In a preferred embodiment of the invention it was an object of the invention to provide for cholesterol lowering or anti-cholestatic compds. It was also an object of the invention to provide for compds. that may be used for the manufacture of anticancer medicaments or apoptosis-inducing medicaments in general.				
IT 499977-78-9	A	499977-78-9		
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
(farnesoid X receptor NR1H4 nuclear receptor binding compds. for treatment of cholesterol or bile acid associated conditions or cancer or to induce apoptosis in relation to gene expression)				
RN 499977-78-9 CAPLUS				
CN Benzenepropanoic acid, 4-[4-[3-[4-(2,6-dichlorophenyl)-5-(1-methylethyl)-5-isoxazolyl]butoxy]- (CA INDEX NAME)				




RN 628332-73-0 CAPLUS

CN Benzenepropanoic acid, 3-[4-[3-[4-(trifluoromethyl)phenyl]-5-isoxazolyl]butoxy]- (CA INDEX NAME)




RN 628332-75-2 CAPLUS

CN Benzenepropanoic acid, 2-ethoxy-4-[4-[3-[4-(trifluoromethyl)phenyl]-5-isoxazolyl]butoxy]- (CA INDEX NAME)

RE.CNT 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L10 ANSWER 21 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN

isoxazolyl]methoxy]- (CA INDEX NAME)



RN 499987-79-0 CAPLUS

CN 2-Propenoic acid, 3-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]- (CA INDEX NAME)

RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L10 ANSWER 22 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN
 AN 2002:964190 CAPLUS Full-text
 DN 138:39272
 TI Preparation of 3-(oxazolylalkoxy)phenyl)propionic acids and analogs as modulators of peroxisome proliferator activated receptors for treatment of diabetes and related conditions
 IN Gosssett, Lynn Stacy; Green, Jonathan Edward; Henry, James Robert; Jones, Winton Dennis, Jr.; Matthews, Donald Paul; Shen, Quan Rong; Smith, Daryl Lynn; Vance, Jennifer Ann; Marshawsky, Alan M.
 PA Eli Lilly and Company, USA
 SO PCT Int. Appl., 438 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002-964190	A1	20020902	WO 2001-485721	20010813

FAN.CMT 1											
PATENT NO.		KIND		DATE		APPLICATION NO.				DATE	
-----		----		----		-----				-----	
PI	EP 581095	A2		19940202		EP 1993-111103				19930712	
R: AT. BR. CH. DE. DK. ES. FR. GB. GR. IE. IT. LI. NL. PT. SE											

CA 2100546	A1	19940125	CA 1993-2100546	19930714
JP 06211748	A	19940802	JP 1993-181305	19930722
AU 9342121	A	19940127	AU 1993-42121	19930723
AU 660226	B2	19930615		
HU 66105	A2	19940928	HU 1993-2150	19930723
ZA 9305332	A	19950123	ZA 1993-5332	19930723

PRA1 DE 1992-4224457
 OS MARPAT 121:8893
 GI



AB The title compds. [I; B = (un)substituted alkyl, C1-4 (un)substituted alkenyl, (un)substituted alkynyl, etc.; R1, R2 = (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, etc.; X, Y = H, halogen, CN, NO2, haloalkyl, alkyl, alkenyl, alkynyl, heteroaryl, heterocyclyl, etc.], useful as agrochem. fungicides, are prepared and I-containing formulations presented. Thus, Me α -(2-hydroxyphenyl)- β -methoxyacrylate was condensed with phenacyl bromide, producing acrylate II, m.p. 76°, which demonstrated 90% inhibitory activity against *Plasmopara viticola* at 250 ppm.


IT 154594-53-1P 154594-53-2P 154594-51-1P
 154594-55-5P 154594-56-6P 154594-69-1P
 154594-73-4P 154594-81-7P 154594-92-0P
 154594-92-1P 154594-94-2P 154594-95-3P
 154594-96-4P 154595-94-7P 154595-95-6P
 154595-96-0P 154595-97-0P

RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of, as agrochem. fungicide)

RN 154594-52-2 CAPLUS


CN Benzenecetic acid, α -(methoxymethylene)-2-[[3-phenyl-5-isoxazolyl)methoxy]-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



RN 154594-53-3 CAPLUS


CN Benzenecetic acid, α -(methoxymethylene)-2-[[3-(2-methylphenyl)-5-isoxazolyl)methoxy]-, methyl ester, (E)- (9CI) (CA INDEX NAME)



RN 154594-69-1 CAPLUS

CN Benzenecetic acid, 2-[[5-(4-chlorophenyl)-3-isoxazolyl)methoxy]- α -(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-70-4 CAPLUS

CN Benzenecetic acid, 2-[[3-(4-chlorophenyl)-5-isoxazolyl)methoxy]- α -(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 154594-81-7 CAPLUS

CN Benzenecetic acid, α -(methoxymethylene)-2-[[3-phenyl-5-isoxazolyl)methoxy]-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.


Double bond geometry as shown.



RN 154594-54-4 CAPLUS

CN Benzenecetic acid, α -(methoxymethylene)-2-[[3-(3-methylphenyl)-5-isoxazolyl)methoxy]-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-55-5 CAPLUS

CN Benzenecetic acid, α -(methoxymethylene)-2-[[3-(4-methylphenyl)-5-isoxazolyl)methoxy]-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-56-6 CAPLUS

CN Benzenecetic acid, 2-[[4,5-dihydro-3-(4-methylphenyl)-5-isoxazolyl)methoxy]- α -(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-92-0 CAPLUS

CN Benzenecetic acid, α -(methoxymethylene)-2-[[5-(4-methylphenyl)-3-isoxazolyl)methoxy]-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-93-1 CAPLUS

CN Benzenecetic acid, α -(methoxymethylene)-2-[[5-(4-methylphenyl)-3-isoxazolyl)methoxy]-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-94-2 CAPLUS

CN Benzenecetic acid, 2-[[4-chloro-5-phenyl-3-isoxazolyl)methoxy]- α -(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-95-3 CAPLUS

CN Benzenecetic acid, 2-[[5-(4-chlorophenyl)-4-methyl-3-isoxazolyl]methoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154594-96-4 CAPLUS

CN Benzenecetic acid, 2-[[5-(3-chlorophenyl)-4-methyl-3-isoxazolyl]methoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154595-04-7 CAPLUS

CN Benzenecetic acid, 2-[[4-chloro-3-(3-fluorophenyl)-5-isoxazolyl]methoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



L10 ANSWER 25 OF 25 CAPLUS COPYRIGHT 2008 ACS ON STN

AN 1990:459160 CAPLUS [Full-text](#)

DN 113:59160

TI Preparation of isoxazol-3-one derivatives as reagents for protecting acids

IN Ito, Takayuki; Nakamura, Takeki

PA Fuji Photo Film Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 17 pp.


CODEN: JXXKXAF

DT Patent


LA Japanese

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
P1 JP 02049776	A	19900220	JP 1988-200603	19880811
PR1 JP 1988-200603		19880811		
OS MARPAT 113:59160				
GI				




AB The title compds. (I; X = halo, sulfonyloxy; R1,R2 = H, substituent; provided that at least one of R1 and R2 is NO₂-substituted aryl or heterocyclyl) are prepared and can be used to protect proton acids having pK_a ≤15, e.g. phenols, carboxylic acids, and sulfonic acids, under mild conditions to form groups stable under weakly basic to acidic conditions, while the selective deprotection is effected by reduction or photochem. reduction under a neutral condition. Thus, a mixture of N-methyl-N-hexadecyl-3-nitro-4-chlorobenzeneulfonamide, 5-tert-butyl-3-hydroxyisoxazole, K₂CO₃, and DMSO was



RN 154595-05-8 CAPLUS

CN Benzenecetic acid, 2-[[4-ethyl-5-phenyl-3-isoxazolyl]methoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 154595-06-9 CAPLUS

CN Benzenecetic acid, 2-[[4-chloro-5-(4-chlorophenyl)-3-isoxazolyl]methoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



RN 154595-07-0 CAPLUS

CN Benzenecetic acid, 2-[[4-chloro-5-(4-methylphenyl)-3-isoxazolyl]methoxy]-α-(methoxymethylene)-, methyl ester, (E)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.

heated 6 h at 60° to give 100% 5-tert-butyl-2-(4-N-methyl-N-hexadecylsulfamoyl-2-nitrophenyl)isoxazolin-3-one which was treated with paraformaldehyde in refluxing AcOH containing ZnCl₂ under a stream of HCl (g) to give 3-oxoisoxazol-4-ylmethyl chloride, i.e., QCl. Versatility, stability, and selectivity of the protecting group O for phenolic OH groups was demonstrated; e.g. treatment of 5-hydroxybenzoxazole derivative (II; R₃ = H) with QCl in refluxing Me₂CO containing K₂CO₃ and KI, hydrolysis of the product II (R₃ = Q) with refluxing 12N aqueous HCl and EtOH to dihydroxyaniline III.HCl (R₄ = R₅ = H, R₆ = Q), and successive acylation of the latter with (Me₃CO₂C)₂O and Ac₂O in pyridine gave III (R₄ = Ac, R₅ = CO₂Me₃, R₆ = Q). Treatment of the latter with trimethylhydroquinone and Et₃N in DMF at 10-25° gave 75% III (R₄ = Ac, R₅ = CO₂Me₃, R₆ = H).


IT 128141-55-9P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, oxoisoxazolylmethyl protecting group in)

RN 128141-55-9 CAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-O-[(5-(1,1-dimethylethyl)-2,3-dihydro-2-[4-[(methyloctadecylamino)sulfonyl]-2-nitrophenyl]-3-oxo-4-isoxazolyl)methyl]- (CA INDEX NAME)

Absolute stereochemistry.



== log hold
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE ENTRY 136.73
TOTAL SESSION 186.43

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)
CA SUBSCRIBER PRICE

SINCE FILE ENTRY -20.00
TOTAL SESSION -20.00

SESSION WILL BE HELD FOR 120 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 11:53:40 ON 16 JAN 2008